

产品名称: Tec Rabbit Monoclonal Antibody
产品货号: AMRe02682



产品概述 (Summary)

产品名称 (Production Name)	Tec Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% Sodium azide and 0.05% protective protein
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TEC
别名 (Alternative Names)	Tyrosine-protein kinase Tec
基因 ID (Gene ID)	7006
蛋白 ID (SwissProt ID)	P42680.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000, ICC/IF 1:50-1:200
蛋白分子量 (Molecular Weight)	Calculated MW: 74 kDa; Observed MW: 58-75 kDa

研究背景 (Background)

Non-receptor tyrosine kinase that contributes to signaling from many receptors and participates as a signal transducer in

产品名称: Tec Rabbit Monoclonal Antibody
产品货号: AMRe02682




multiple downstream pathways, including regulation of the actin cytoskeleton. Plays a redundant role to ITK in regulation of the adaptive immune response. Regulates the development, function and differentiation of conventional T-cells and nonconventional NKT-cells. Required for TCR-dependent IL2 gene induction. Phosphorylates DOK1, one CD28-specific substrate, and contributes to CD28-signaling. Mediates signals that negatively regulate IL2RA expression induced by TCR cross-linking. Plays a redundant role to BTK in BCR-signaling for B-cell development and activation, especially by phosphorylating STAP1, a BCR-signaling protein. Required in mast cells for efficient cytokine production. Involved in both growth and differentiation mechanisms of myeloid cells through activation by the granulocyte colony-stimulating factor CSF3, a critical cytokine to promoting the growth, differentiation, and functional activation of myeloid cells. Participates in platelet signaling downstream of integrin activation. Cooperates with JAK2 through reciprocal phosphorylation to mediate cytokine-driven activation of FOS transcription. GRB10, a negative modifier of the FOS activation pathway, is another substrate of TEC. TEC is involved in G protein-coupled receptor- and integrin-mediated signalings in blood platelets. Plays a role in hepatocyte proliferation and liver regeneration and is involved in HGF-induced ERK signaling pathway. TEC regulates also FGF2 unconventional secretion (endoplasmic reticulum (ER)/Golgi-independent mechanism) under various physiological conditions through phosphorylation of FGF2 'Tyr-215'. May also be involved in the regulation of osteoclast differentiation.

研究领域 (Research Area)


Signal Transduction

图片 (Image Data)



Western blot analysis of Tec in K562 lysates using Tec antibody.

产品名称: Tec Rabbit Monoclonal Antibody
产品货号: AMRe02682



Immunocytochemistry analysis of Tec(green) in Hela using Tec antibody,and DAPI(blue)

注意事项 (Note)

For research use only .