

产品名称: Phospho-ATF2 (T71) (7E18) Rabbit Monoclonal Antibody
产品货号: AMRe05855



产品概述 (Summary)

产品名称 (Production Name)	Phospho-ATF2 (T71) (7E18) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF, IP
种属反应性 (Reactivity)	Human, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% protective protein.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ATF2
别名 (Alternative Names)	ATF2, Activating 2, CREB2, CREBP1, Cyclic-AMP-dependent ATF-2, HB16, MXBP protein, cAMP response element binding protein CRE-BP1;
基因 ID (Gene ID)	1386.0
蛋白 ID (SwissProt ID)	P15336.A synthetic phosphopeptide corresponding to residues surrounding Thr71 of human ATF2

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000, ICC/IF 1:200-1:500, IP 1:20-1:50
蛋白分子量 (Molecular Weight)	55kDa

产品名称: Phospho-ATF2 (T71) (7E18) Rabbit Monoclonal Antibody
产品货号: AMRe05855


研究背景 (Background)

The transcription factor ATF-2 (also called CRE-BP1) binds to both AP-1 and CRE DNA response elements and is a member of the ATF/CREB family of leucine zipper proteins. ATF-2 interacts with a variety of viral oncoproteins and cellular tumor suppressors and is a target of the SAPK/JNK and p38 MAP kinase signaling pathways. Various forms of cellular stress, including genotoxic agents, inflammatory cytokines, and UV irradiation, stimulate the transcriptional activity of ATF-2. Cellular stress activates ATF-2 by phosphorylation of Thr69 and Thr71. Transcriptional activator which regulates the transcription of various genes, including those involved in anti-apoptosis, cell growth, and DNA damage response. Dependent on its binding partner, binds to CRE (cAMP response element) consensus sequences (5'-TGACGTCA- 3') or to AP-1 (activator protein 1) consensus sequences (5'-TGACTCA- 3'). In the nucleus, contributes to global transcription and the DNA damage response, in addition to specific transcriptional activities that are related to cell development, proliferation and death. In the cytoplasm, interacts with and perturbs HK1- and VDAC1-containing complexes at the mitochondrial outer membrane, thereby impairing mitochondrial membrane potential, inducing mitochondrial leakage and promoting cell death. The phosphorylated form (mediated by ATM) plays a role in the DNA damage response and is involved in the ionizing radiation (IR)-induced S phase checkpoint control and in the recruitment of the MRN complex into the IR-induced foci (IRIF). Exhibits histone acetyltransferase (HAT) activity which specifically acetylates histones H2B and H4 in vitro (PubMed:10821277). In concert with CUL3 and RBX1, promotes the degradation of KAT5 thereby attenuating its ability to acetylate and activate ATM. Can elicit oncogenic or tumor suppressor activities depending on the tissue or cell type.

研究领域 (Research Area)

Epigenetics and Nuclear Signaling

图片 (Image Data)



Western blot detection of Phospho-ATF2 (Thr71) in 3T3, HeLa cell lysates using Phospho-ATF2 (Thr71) antibody (1:1000 diluted).

产品名称: Phospho-ATF2 (T71) (7E18) Rabbit Monoclonal
Antibody
产品货号: AMRe05855



注意事项 (Note)

For research use only .