

产品货号: AMRe05906

产品概述 (Summary)

产品名称 (Production Name) Phospho-FAK (Y397) (10H3) Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit 应用 (Application) WB,ICC/IF

种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) lgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New 储存溶液 (Buffer)

type preservative N and 0.05% protective protein.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) PTK2

EC 2.7.10.2; FADK 1; FAK1; Focal adhesion kinase 1; PTK2; Protein- tyrosine 別名 (Alternative Names)

kinase 2; pp125FAK;

基因 ID (Gene ID) 5747.0

Q05397.A synthetic phosphopeptide corresponding to residues surrounding 蛋白 ID(SwissProt ID)

Tyr397 of human FAK

产品应用(Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ICC/IF 1:100-1:200

蛋白分子量 (Molecular Weight) 119kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

Antibody

产品货号: AMRe05906



研究背景 (Background)

This gene encodes a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. Non-receptor protein-tyrosine kinase that plays an essential role in regulating cell migration, adhesion, spreading, reorganization of the actin cytoskeleton, formation and disassembly of focal adhesions and cell protrusions, cell cycle progression, cell proliferation and apoptosis. Required for early embryonic development and placenta development. Required for embryonic angiogenesis, normal cardiomyocyte migration and proliferation, and normal heart development. Regulates axon growth and neuronal cell migration, axon branching and synapse formation; required for normal development of the nervous system. Plays a role in osteogenesis and differentiation of osteoblasts. Functions in integrin signal transduction, but also in signaling downstream of numerous growth factor receptors, G-protein coupled receptors (GPCR), EPHA2, netrin receptors and LDL receptors. Forms multisubunit signaling complexes with SRC and SRC family members upon activation; this leads to the phosphorylation of additional tyrosine residues, creating binding sites for scaffold proteins, effectors and substrates. Regulates numerous signaling pathways. Promotes activation of phosphatidylinositol 3-kinase and the AKT1 signaling cascade. Promotes activation of MAPK1/ERK2, MAPK3/ERK1 and the MAP kinase signaling cascade. Promotes localized and transient activation of quanine nucleotide exchange factors (GEFs) and GTPase-activating proteins (GAPs), and thereby modulates the activity of Rho family GTPases. Signaling via CAS family members mediates activation of RAC1. Recruits the ubiquitin ligase MDM2 to P53/TP53 in the nucleus, and thereby regulates P53/TP53 activity, P53/TP53 ubiquitination and proteasomal degradation. Phosphorylates SRC; this increases SRC kinase activity. Phosphorylates ACTN1, ARHGEF7, GRB7, RET and WASL. Promotes phosphorylation of PXN and STAT1; most likely PXN and STAT1 are phosphorylated by a SRC family kinase that is recruited to autophosphorylated PTK2/FAK1, rather than by PTK2/FAK1 itself. Promotes phosphorylation of BCAR1; GIT2 and SHC1; this requires both SRC and PTK2/FAK1. Promotes phosphorylation of BMX and PIK3R1. Isoform 6 (FRNK) does not contain a kinase domain and inhibits PTK2/FAK1 phosphorylation and signaling. Its enhanced expression can attenuate the nuclear accumulation of LPXN and limit its ability to enhance serum response factor (SRF)-dependent gene transcription.

研究领域(Research Area)


Cardiovascular

图片 (Image Data)

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838



产品货号: AMRe05906



Western blot detection of Phospho-FAK (Tyr397) in C6 cell lysates using Phospho-FAK (Tyr397) antibody(1:1000 diluted).

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838