

产品名称: ACVRL1 (9G1) Rabbit Monoclonal Antibody
产品货号: AMRe06570



产品概述 (Summary)

产品名称 (Production Name)	ACVRL1 (9G1) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ACVRL1
别名 (Alternative Names)	SKR3; Activin receptor-like kinase 1; ALK-1; TGF-B superfamily receptor type I; TSR-I; ACVRL1; ACVRLK1; ALK1;
基因 ID (Gene ID)	94.0
蛋白 ID (SwissProt ID)	P37023.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000
蛋白分子量 (Molecular Weight)	56kDa

产品名称: ACVRL1 (9G1) Rabbit Monoclonal Antibody
产品货号: AMRe06570




研究背景 (Background)

On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. Receptor for TGF-beta. May bind activin as well. Type I receptor for TGF-beta family ligands BMP9/GDF2 and BMP10 and important regulator of normal blood vessel development. On ligand binding, forms a receptor complex consisting of two type II and two type I transmembrane serine/threonine kinases. Type II receptors phosphorylate and activate type I receptors which autophosphorylate, then bind and activate SMAD transcriptional regulators. May bind activin as well.

研究领域 (Research Area)

Cytokine-cytokine receptor interaction;TGF-beta;

图片 (Image Data)



Western blot analysis of ACVRL1 expression in (1) Jurkat cell lysate; (2) Mouse kidney lysate; (3) Rat kidney lysate.

注意事项 (Note)

For research use only .