

产品名称: Bcl10 (9M3) Rabbit Monoclonal Antibody
产品货号: AMRe07497



产品概述 (Summary)

产品名称 (Production Name)	Bcl10 (9M3) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,FC
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	BCL10
别名 (Alternative Names)	BCL10; CARMEN; CIPER; CLAP; c-E10; mE10;
基因 ID (Gene ID)	8915.0
蛋白 ID (SwissProt ID)	O95999.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000,FC 1:200-1:1000
蛋白分子量 (Molecular Weight)	26kDa

研究背景 (Background)

产品名称: Bcl10 (9M3) Rabbit Monoclonal Antibody
产品货号: AMRe07497




Promotes apoptosis, pro-caspase-9 maturation and activation of NF-kappa-B via NIK and IKK. May be an adapter protein between upstream TNFR1-TRADD-RIP complex and the downstream NIK-IKK-IKAP complex. Is a substrate for MALT1. Plays a key role in both adaptive and innate immune signaling by bridging CARD domain-containing proteins to immune activation (PubMed: [10187770](http://www.uniprot.org/citations/10187770), PubMed: [10364242](http://www.uniprot.org/citations/10364242), PubMed: [10400625](http://www.uniprot.org/citations/10400625), PubMed: [25365219](http://www.uniprot.org/citations/25365219), PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Acts by channeling adaptive and innate immune signaling downstream of CARD domain-containing proteins CARD9, CARD11 and CARD14 to activate NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Recruited by activated CARD domain-containing proteins: homooligomerized CARD domain-containing proteins form a nucleating helical template that recruits BCL10 via CARD-CARD interaction, thereby promoting polymerization of BCL10, subsequent recruitment of MALT1 and formation of a CBM complex (PubMed: [24074955](http://www.uniprot.org/citations/24074955)). This leads to activation of NF-kappa-B and MAP kinase p38 (MAPK11, MAPK12, MAPK13 and/or MAPK14) pathways which stimulate expression of genes encoding pro-inflammatory cytokines and chemokines (PubMed: [18287044](http://www.uniprot.org/citations/18287044), PubMed: [27777308](http://www.uniprot.org/citations/27777308), PubMed: [24074955](http://www.uniprot.org/citations/24074955), PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Activated by CARD9 downstream of C-type lectin receptors; CARD9-mediated signals are essential for antifungal immunity (PubMed: [26488816](http://www.uniprot.org/citations/26488816)). Activated by CARD11 downstream of T-cell receptor (TCR) and B-cell receptor (BCR) (PubMed: [18264101](http://www.uniprot.org/citations/18264101), PubMed: [18287044](http://www.uniprot.org/citations/18287044), PubMed: [27777308](http://www.uniprot.org/citations/27777308), PubMed: [24074955](http://www.uniprot.org/citations/24074955)). Promotes apoptosis, pro-caspase-9 maturation and activation of NF-kappa-B via NIK and IKK (PubMed: [10187815](http://www.uniprot.org/citations/10187815)).

研究领域 (Research Area)

Cell Biology

图片 (Image Data)

产品名称: Bcl10 (9M3) Rabbit Monoclonal Antibody
产品货号: AMRe07497



Western blot analysis of Bcl10 expression in HeLa cell lysate.

注意事项 (Note)

For research use only .