

产品名称: CD11b (6H15) Rabbit Monoclonal Antibody
产品货号: AMRe08195



产品概述 (Summary)

产品名称 (Production Name)	CD11b (6H15) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,IF-P
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ITGAM
别名 (Alternative Names)	CR3A; MO1A; CD11B; MAC-1; MAC1A; SLEB6;ITGAM
基因 ID (Gene ID)	3684.0
蛋白 ID (SwissProt ID)	P11215.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:500-1:2000,IF-P 1:500-1:2000
蛋白分子量 (Molecular Weight)	127kDa

研究背景 (Background)

产品名称: CD11b (6H15) Rabbit Monoclonal Antibody
产品货号: AMRe08195




ITGAM, also named as CD11B and CR3A, belongs to the integrin alpha chain family. It is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles. ITGAM is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin ITGAM/ITGB2 is implicated in various adhesive interactions of monocytes, macrophages and granulocytes as well as in mediating the uptake of complement-coated particles and pathogens (PubMed: [9558116](http://www.uniprot.org/citations/9558116), PubMed: [20008295](http://www.uniprot.org/citations/20008295)). It is identical with CR-3, the receptor for the iC3b fragment of the third complement component. It probably recognizes the R-G-D peptide in C3b. Integrin ITGAM/ITGB2 is also a receptor for fibrinogen, factor X and ICAM1. It recognizes P1 and P2 peptides of fibrinogen gamma chain. Regulates neutrophil migration (PubMed: [28807980](http://www.uniprot.org/citations/28807980)). In association with beta subunit ITGB2/CD18, required for CD177-PRTN3-mediated activation of TNF primed neutrophils (PubMed: [21193407](http://www.uniprot.org/citations/21193407)). May regulate phagocytosis-induced apoptosis in extravasated neutrophils (By similarity). May play a role in mast cell development (By similarity). Required with TYROBP/DAP12 in microglia to control production of microglial superoxide ions which promote the neuronal apoptosis that occurs during brain development (By similarity).

研究领域 (Research Area)

Cell adhesion molecules (CAMs); Hematopoietic cell lineage; Leukocyte transendothelial migration; Regulates Actin and Cytoskeleton;

图片 (Image Data)



Western blot analysis of CD11b expression in RAW264.7 cell lysate.

注意事项 (Note)

For research use only .