

产品名称: Cdc25C (11W11) Rabbit Monoclonal Antibody
产品货号: AMRe08507



产品概述 (Summary)

产品名称 (Production Name)	Cdc25C (11W11) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CDC25C
别名 (Alternative Names)	CDC25; CDC25C; Cell division cycle 25 homolog C; MP1P3; PPP1R60;
基因 ID (Gene ID)	995.0
蛋白 ID (SwissProt ID)	P30307.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000
蛋白分子量 (Molecular Weight)	53kDa

研究背景 (Background)

产品名称: Cdc25C (11W11) Rabbit Monoclonal Antibody
产品货号: AMRe08507




Functions as a dosage-dependent inducer in mitotic control. It is a tyrosine protein phosphatase required for progression of the cell cycle. It directly dephosphorylates CDK1 and activate its kinase activity. Functions as a dosage-dependent inducer in mitotic control. Tyrosine protein phosphatase required for progression of the cell cycle. When phosphorylated, highly effective in activating G2 cells into prophase. Directly dephosphorylates CDK1 and activates its kinase activity.

研究领域 (Research Area)

Cell Biology

图片 (Image Data)



Western blot analysis of Cdc25C expression in K562 cell lysate.

注意事项 (Note)

For research use only .