

产品名称: CDKN2A/p14ARF (18X15) Rabbit Monoclonal Antibody
产品货号: AMRe08578



产品概述 (Summary)

产品名称 (Production Name)	CDKN2A/p14ARF (18X15) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF, FC, IP
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CDKN2A
别名 (Alternative Names)	ARF; CDK4I; CDKN2; CDKN2A; CMM2; INK4; INK4a; MLM; MTS1; p14; p16; p16INK4a; P19ARF; TP16;
基因 ID (Gene ID)	1029.0
蛋白 ID (SwissProt ID)	Q8N726.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000, ICC/IF 1:100-1:200, FC 1:50-1:200, IP 1:50-1:100
蛋白分子量 (Molecular Weight)	14kDa

产品名称: CDKN2A/p14ARF (18X15) Rabbit Monoclonal Antibody
产品货号: AMRe08578




研究背景 (Background)

The gene for CDKN2A generates several transcripts/proteins which differ from each other in their first exons. Three of these transcripts are generated by alternative splicing (isoform 1 a.k.a p16INK4A, isoform 2 and isoform 3 a.k.a p12), two of which are known to function as inhibitors of CDK4 kinase. Capable of inducing cell cycle arrest in G1 and G2 phases. Acts as a tumor suppressor. Binds to MDM2 and blocks its nucleocytoplasmic shuttling by sequestering it in the nucleolus. This inhibits the oncogenic action of MDM2 by blocking MDM2-induced degradation of p53 and enhancing p53-dependent transactivation and apoptosis. Also induces G2 arrest and apoptosis in a p53-independent manner by preventing the activation of cyclin B1/CDC2 complexes. Binds to BCL6 and down-regulates BCL6-induced transcriptional repression. Binds to E2F1 and MYC and blocks their transcriptional activator activity but has no effect on MYC transcriptional repression. Binds to TOP1/TOPOI and stimulates its activity. This complex binds to rRNA gene promoters and may play a role in rRNA transcription and/or maturation. Interacts with NPM1/B23 and promotes its polyubiquitination and degradation, thus inhibiting rRNA processing. Interacts with COMMD1 and promotes its 'Lys63'-linked polyubiquitination. Interacts with UBE2I/UBC9 and enhances sumoylation of a number of its binding partners including MDM2 and E2F1. Binds to HUWE1 and represses its ubiquitin ligase activity. May play a role in controlling cell proliferation and apoptosis during mammary gland development. Isoform smARF may be involved in regulation of autophagy and caspase-independent cell death; the short-lived mitochondrial isoform is stabilized by C1QBP.

研究领域 (Research Area)

Stem cell pathway; Cell Cycle

图片 (Image Data)



Western blot analysis of CDKN2A/p14ARF expression in PC3 cell lysate.

注意事项 (Note)

For research use only .