

产品货号: AMRe08639



产品概述 (Summary)

产品名称 (Production Name) CENPE (15A19) Rabbit Monoclonal Antibody

描述 (**Description**) Recombinant rabbit monoclonal antibody

宿主 (Host)Rabbit应用 (Application)WB种属反应性 (Reactivity)Human

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% New

储存溶液 (Buffer) type preservative N and 50% glycerol. Store at +4°C short term. Store at -

20°C long term. Avoid freeze / thaw cycle.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CENPE

CENP E; Centromere associated protein E; Centromere protein E 312kDa; 別名 (Alternative Names)

KIF10; Kinesin family member 10; Kinesin related protein; PPP1R61;

基因 ID (Gene ID) 1062.0 蛋白 ID (SwissProt ID) Q02224.

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:1000-1:5000

蛋白分子量 (Molecular Weight) 316kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: CENPE (15A19) Rabbit Monoclonal Antibody

产品货号: AMRe08639



研究背景 (Background)

CENPE is a 250-300 kDa human centromere-associated kinesin-like motor protein that accumulates in G2 phase. Microtubule plus-end-directed kinetochore motor which plays an important role in chromosome congression, microtubulekinetochore conjugation and spindle assembly checkpoint activation. Drives chromosome congression (alignment of chromosomes at the spindle equator resulting in the formation of the metaphase plate) by mediating the lateral sliding of polar chromosomes along spindle microtubules towards the spindle equator and by aiding the establishment and maintenance of connections between kinetochores and spindle microtubules (PubMed: 7889940, PubMed:23891108, PubMed: 25395579). The transport of pole-proximal chromosomes towards the spindle equator is favored by microtubule tracks that are detyrosinated (PubMed: 25908662). Acts as a processive bi-directional tracker of dynamic microtubule tips; after chromosomes have congressed, continues to play an active role at kinetochores, enhancing their links with dynamic microtubule ends (PubMed: 23955301). Suppresses chromosome congression in NDC80-depleted cells and contributes positively to congression only when microtubules are stabilized (PubMed: 25743205). Plays an important role in the formation of stable attachments between kinetochores and spindle microtubules (PubMed: 17535814) The stabilization of kinetochore- microtubule attachment also requires CENPE-dependent localization of other proteins to the kinetochore including BUB1B, MAD1 and MAD2. Plays a role in spindle assembly checkpoint activation (SAC) via its interaction with BUB1B resulting in the activation of its kinase activity, which is important for activating SAC. Necessary for the mitotic checkpoint signal at individual kinetochores to prevent aneuploidy due to single chromosome loss (By similarity).

研究领域(Research Area)


Epigenetics and Nuclear Signaling

图片 (Image Data)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品货号: AMRe08639





Western blot analysis of CENPE expression in HepG2 cell lysate.

注意事项 (Note)

For research use only .

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838