

产品名称: Emi1 (16Z12) Rabbit Monoclonal Antibody
产品货号: AMRe10439



产品概述 (Summary)

产品名称 (Production Name)	Emi1 (16Z12) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	FBXO5
别名 (Alternative Names)	EMI1; FBX5; Fbxo31; fbxo5;
基因 ID (Gene ID)	26271.0
蛋白 ID (SwissProt ID)	Q9UKT4.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000
蛋白分子量 (Molecular Weight)	50kDa

研究背景 (Background)

产品名称: Emi1 (16Z12) Rabbit Monoclonal Antibody
产品货号: AMRe10439



Regulates progression through early mitosis by inhibiting the anaphase promoting complex/cyclosome (APC). Binds to the APC activators CDC20 and FZR1/CDH1 to prevent APC activation. Can also bind directly to the APC to inhibit substrate-binding. Regulator of APC activity during mitotic and meiotic cell cycle (PubMed:17485488, PubMed:17234884, PubMed:17875940, PubMed:23708001, PubMed:23708605, PubMed:16921029). During mitotic cell cycle plays a role as both substrate and inhibitor of APC-FZR1 complex (PubMed:29875408, PubMed:17485488, PubMed:17234884, PubMed:17875940, PubMed:23708001, PubMed:23708605, PubMed:16921029). During G1 phase, plays a role as substrate of APC-FZR1 complex E3 ligase (PubMed:29875408). Then switches as an inhibitor of APC-FZR1 complex during S and G2 leading to cell-cycle commitment (PubMed:29875408). As APC inhibitor, prevents the degradation of APC substrates at multiple levels: by interacting with APC and blocking access of APC substrates to the D-box coreceptor, formed by FZR1 and ANAPC10; by suppressing ubiquitin ligation and chain elongation by APC by preventing the UBE2C and UBE2S activities (PubMed:23708605, PubMed:23708001, PubMed:16921029). Plays a role in genome integrity preservation by coordinating DNA replication with mitosis through APC inhibition in interphase to stabilize CCNA2 and GMNN in order to promote mitosis and prevent rereplication and DNA damage-induced cellular senescence (PubMed:17234884, PubMed:17485488, PubMed:17875940). During oocyte maturation, plays a role in meiosis through inactivation of APC-FZR1 complex. Inhibits APC through RPS6KA2 interaction that increases FBXO5 affinity for CDC20 leading to the metaphase arrest of the second meiotic division before fertilization (By similarity). Controls entry into the first meiotic division through inactivation of APC-FZR1 complex (By similarity). Promotes migration and osteogenic differentiation of mesenchymal stem cells (PubMed:29850565).


产品名称: Emi1 (16Z12) Rabbit Monoclonal Antibody
产品货号: AMRe10439



研究领域 (Research Area)

Cell Biology

图片 (Image Data)



Western blot analysis of Emi1 expression in HepG2 cell lysate.

注意事项 (Note)

For research use only .