

产品名称: FANCD2 (16Q3) Rabbit Monoclonal Antibody
产品货号: AMRe10826



产品概述 (Summary)

产品名称 (Production Name)	FANCD2 (16Q3) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,FC,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	FANCD2
别名 (Alternative Names)	FA D2; FA4; FAC D2; FACD 2; FACD; FACD2; FAD; FAD2; FANCD 2; FANCD; FANCD2; Fanconi anemia group D2 protein; FLJ23826; Protein FACD2; Type 4 Fanconi pancytopenia;
基因 ID (Gene ID)	2177.0
蛋白 ID (SwissProt ID)	Q9BXW9.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000,IHC 1:50-1:200,ICC/IF 1:200-1:500,FC 1:100-1:200,IP 1:10-1:100
----------------------	---

产品名称: FANCD2 (16Q3) Rabbit Monoclonal Antibody
产品货号: AMRe10826




蛋白分子量 (Molecular Weight) 164kDa

研究背景 (Background)

Required for maintenance of chromosomal stability. Promotes accurate and efficient pairing of homologs during meiosis. Involved in the repair of DNA double-strand breaks, both by homologous recombination and single-strand annealing. Required for maintenance of chromosomal stability. Promotes accurate and efficient pairing of homologs during meiosis. Involved in the repair of DNA double-strand breaks, both by homologous recombination and single-strand annealing. May participate in S phase and G2 phase checkpoint activation upon DNA damage. Plays a role in preventing breakage and loss of missegregating chromatin at the end of cell division, particularly after replication stress. Required for the targeting, or stabilization, of BLM to non-centromeric abnormal structures induced by replicative stress. Promotes BRCA2/FANCD1 loading onto damaged chromatin. May also be involved in B-cell immunoglobulin isotype switching.

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of FANCD2 expression in HeLa cell lysate.

注意事项 (Note)

For research use only .