产品名称: GluR1 (12Y10) Rabbit Monoclonal Antibody

产品货号: AMRe11490



产品概述 (Summary)

产品名称 (Production Name) GluR1 (12Y10) Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit 应用 (Application) WB,IP

种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% New

储存溶液 (Buffer) type preservative N and 50% glycerol. Store at +4°C short term. Store at -

20°C long term. Avoid freeze / thaw cycle.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) GRIA1

Glutamate receptor 1; GluR-1; AMPA-selective glutamate receptor 1; GluR-A;

别名 (Alternative Names) GluR-K1; GluRA; GluRK1; Glutamate receptor ionotropic, AMPA 1; GluA1;

GRIA1; GLUH1; GLUR1;

基因 ID (Gene ID) 2890.0 **蛋白 ID (SwissProt ID)** P42261.

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IP 1:10-1:100

蛋白分子量 (Molecular Weight) 102kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: GluR1 (12Y10) Rabbit Monoclonal Antibody

产品货号: AMRe11490




研究背景 (Background)

AMPA- (α-amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid), kainate-, and NMDA- (N-methyl-D-aspartate) receptors are the three main families of ionotropic glutamate-gated ion channels. AMPA receptors (AMPARs) are comprised of four subunits (GluR 1-4), which assemble as homo- or hetero-tetramers to mediate the majority of fast excitatory transmissions in the central nervous system. AMPARs are implicated in synapse formation, stabilization, and plasticity. Ionotropic glutamate receptor. L-glutamate acts as an excitatory neurotransmitter at many synapses in the central nervous system. Binding of the excitatory neurotransmitter L-glutamate induces a conformation change, leading to the opening of the cation channel, and thereby converts the chemical signal to an electrical impulse. The receptor then desensitizes rapidly and enters a transient inactive state, characterized by the presence of bound agonist. In the presence of CACNG4 or CACNG7 or CACNG8, shows resensitization which is characterized by a delayed accumulation of current flux upon continued application of glutamate.

研究领域 (Research Area)

Neuroscience

图片 (Image Data)



Western blot analysis of GluR1 expression in Human brain lysate.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838