

产品名称: JAB1 (7S6) Rabbit Monoclonal Antibody
产品货号: AMRe12808



产品概述 (Summary)

产品名称 (Production Name)	JAB1 (7S6) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% protective protein.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	COPS5
别名 (Alternative Names)	cops5; CSN 5; CSN5; JAB 1; MGC3149; MOV34; SGN5;
基因 ID (Gene ID)	10987.0
蛋白 ID (SwissProt ID)	Q92905.A synthetic peptide of human JAB1

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000
蛋白分子量 (Molecular Weight)	38kDa

研究背景 (Background)

Probable protease subunit of the COP9 signalosome complex (CSN), a complex involved in various cellular and


产品名称: JAB1 (7S6) Rabbit Monoclonal Antibody
产品货号: AMRe12808



developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of the SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. Probable protease subunit of the COP9 signalosome complex (CSN), a complex involved in various cellular and developmental processes. The CSN complex is an essential regulator of the ubiquitin (Ubl) conjugation pathway by mediating the deneddylation of the cullin subunits of the SCF-type E3 ligase complexes, leading to decrease the Ubl ligase activity of SCF-type complexes such as SCF, CSA or DDB2. The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN, IkappaBalpha/NFKBIA, ITPK1 and IRF8, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects degradation by the Ubl system, respectively. In the complex, it probably acts as the catalytic center that mediates the cleavage of Nedd8 from cullins. It however has no metalloprotease activity by itself and requires the other subunits of the CSN complex. Interacts directly with a large number of proteins that are regulated by the CSN complex, confirming a key role in the complex. Promotes the proteasomal degradation of BRSK2.

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from PC-12 cells using RM5724 at 1:1000.

注意事项 (Note)

For research use only .