产品名称: MCM2 (6N7) Rabbit Monoclonal Antibody

产品货号: AMRe13718



## 产品概述 (Summary)

产品名称 (Production Name) MCM2 (6N7) Rabbit Monoclonal Antibody

描述 (**Description**) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit

**应用 (Application)** WB,IHC,ICC/IF,FC,IP

种属反应性 (Reactivity) Human

## 产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% New

储存溶液 (Buffer) type preservative N and 50% glycerol. Store at +4°C short term. Store at -

20°C long term. Avoid freeze / thaw cycle.

纯化方式 (Purification) Affinity purification

## 免疫原信息 (Immunogen)

基因名 (Gene Name) MCM2

别名 (Alternative Names) BM28; CCNL1; cdc19; CDCL1; Cyclin like 1; MCM2;

**基因 ID (Gene ID)** 4171.0 **蛋白 ID (SwissProt ID)** P49736.

# 产品应用(Application)

**稀释比 (Dilution Ratio)** WB 1:500-1:2000,IHC 1:20-1:100,ICC/IF 1:20-1:50,FC 1:20-1:50,IP 1:20-1:50

蛋白分子量 (Molecular Weight) 102kDa

## 研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838



产品货号: AMRe13718




Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Acts as component of the MCM2-7 complex (MCM complex) which is the putative replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for the entry in S phase and for cell division. Plays a role in terminally differentiated hair cells development of the cochlea and induces cells apoptosis.

## 研究领域 (Research Area)

**Epigenetics and Nuclear Signaling** 

## 图片 (Image Data)



Western blot analysis of MCM2 expression in HeLa cell lysate.

#### 注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838