

产品名称: MCM5 (1M15) Rabbit Monoclonal Antibody
产品货号: AMRe13723



产品概述 (Summary)

产品名称 (Production Name)	MCM5 (1M15) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,FC,IP
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% protective protein.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MCM5
别名 (Alternative Names)	CDC46; MCM5; MGC5315; P1 CDC46;
基因 ID (Gene ID)	4174.0
蛋白 ID (SwissProt ID)	P33992.A synthetic peptide of human MCM5

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:2000,IHC 1:50-1:100,ICC/IF 1:20-1:50,FC 1:20-1:50,IP 1:20-1:50
蛋白分子量 (Molecular Weight)	82kDa

研究背景 (Background)

MCM5 Belongs to the MCM family. Acts as component of the MCM2-7 complex (MCM complex) which is the putative

产品名称: MCM5 (1M15) Rabbit Monoclonal Antibody
产品货号: AMRe13723




replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity.

研究领域 (Research Area)

Epigenetics and Nuclear Signaling

图片 (Image Data)



Western blot detection of MCM5 in Jurkat,C6,HeLa cell lysates using MCM5 antibody(1:1000 diluted).

注意事项 (Note)

For research use only .