

产品名称: MMP2 (12D11) Rabbit Monoclonal Antibody
产品货号: AMRe13986



产品概述 (Summary)

产品名称 (Production Name)	MMP2 (12D11) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF, FC
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% New type preservative N and 0.05% protective protein.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MMP2
别名 (Alternative Names)	CLG 4; CLG 4A; CLG4; CLG4A; Gelatinase A; Gelatinase alpha; Gelatinase neutrophil; MMP 2; MMP II; MMP2; MONA; PEX; TBE 1;
基因 ID (Gene ID)	4313.0
蛋白 ID (SwissProt ID)	P08253.A synthetic peptide of human MMP2

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000, ICC/IF 1:100-1:200, FC 1:200-1:500
蛋白分子量 (Molecular Weight)	74kDa

研究背景 (Background)

产品名称: MMP2 (12D11) Rabbit Monoclonal Antibody
产品货号: AMRe13986




Ubiquitous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta-type CGRP promoting vasoconstriction. Ubiquitous metalloproteinase that is involved in diverse functions such as remodeling of the vasculature, angiogenesis, tissue repair, tumor invasion, inflammation, and atherosclerotic plaque rupture. As well as degrading extracellular matrix proteins, can also act on several nonmatrix proteins such as big endothelial 1 and beta- type CGRP promoting vasoconstriction. Also cleaves KISS at a Gly-|-Leu bond. Appears to have a role in myocardial cell death pathways. Contributes to myocardial oxidative stress by regulating the activity of GSK3beta. Cleaves GSK3beta in vitro. Involved in the formation of the fibrovascular tissues in association with MMP14. [Isoform 2]: Mediates the proteolysis of CHUK/IKKA and initiates a primary innate immune response by inducing mitochondrial- nuclear stress signaling with activation of the pro-inflammatory NF- kappaB, NFAT and IRF transcriptional pathways.

研究领域 (Research Area)

Cardiovascular

图片 (Image Data)



Western blot detection of MMP2 in mouse lung, mouse heart cell lysates using MMP2 antibody (1:1000 diluted).

注意事项 (Note)

For research use only .