

产品名称: NR2C2 / TR4 (9S17) Rabbit Monoclonal Antibody
产品货号: AMRe14876



产品概述 (Summary)

产品名称 (Production Name)	NR2C2 / TR4 (9S17) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NR2C2
别名 (Alternative Names)	hTAK1; Nr2c2; Nuclear hormone receptor TR4; Nuclear receptor subfamily 2 group C member 2; Orphan nuclear receptor TAK1; Orphan nuclear receptor TR4; TAK1; Testicular nuclear receptor 4; Testicular receptor 4; TR2R1; TR4; TR4 nuclear hormone receptor;
基因 ID (Gene ID)	7182.0
蛋白 ID (SwissProt ID)	P49116.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000,IP 1:10-1:100
----------------------	--------------------------------

产品名称: NR2C2 / TR4 (9S17) Rabbit Monoclonal Antibody
产品货号: AMRe14876




蛋白分子量 (Molecular Weight) 65kDa

研究背景 (Background)

Orphan nuclear receptor that can act as a repressor or activator of transcription. An important repressor of nuclear receptor signaling pathways such as retinoic acid receptor, retinoid X, vitamin D3 receptor, thyroid hormone receptor and estrogen receptor pathways. May regulate gene expression during the late phase of spermatogenesis. Orphan nuclear receptor that can act as a repressor or activator of transcription. An important repressor of nuclear receptor signaling pathways such as retinoic acid receptor, retinoid X, vitamin D3 receptor, thyroid hormone receptor and estrogen receptor pathways. May regulate gene expression during the late phase of spermatogenesis. Together with NR2C1, forms the core of the DRED (direct repeat erythroid-definitive) complex that represses embryonic and fetal globin transcription including that of GATA1. Binds to hormone response elements (HREs) consisting of two 5'-AGGTCA-3' half site direct repeat consensus sequences. Plays a fundamental role in early embryonic development and embryonic stem cells. Required for normal spermatogenesis and cerebellum development. Appears to be important for neurodevelopmentally regulated behavior (By similarity). Activates transcriptional activity of LHCG. Antagonist of PPARA-mediated transactivation.

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of NR2C2/TR4 expression in PC-3 cell lysate.

注意事项 (Note)

For research use only .