

产品名称: PAK3 (1O10) Rabbit Monoclonal Antibody
产品货号: AMRe15708



产品概述 (Summary)

产品名称 (Production Name)	PAK3 (1O10) Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% New type preservative N and 50% glycerol. Store at +4°C short term. Store at -20°C long term. Avoid freeze / thaw cycle.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PAK3
别名 (Alternative Names)	Beta PAK; bPAK; CDKN1A; hPAK3; MRX30; MRX47; OPHN3; PAK3beta; Pak65alpha; Pak65beta; Stk4;
基因 ID (Gene ID)	5063.0
蛋白 ID (SwissProt ID)	O75914.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:2000-1:20000,IHC 1:100-1:200
蛋白分子量 (Molecular Weight)	62kDa

产品名称: PAK3 (1010) Rabbit Monoclonal Antibody
产品货号: AMRe15708




研究背景 (Background)

PAK proteins are critical effectors that link Rho GTPases to cytoskeleton reorganization and nuclear signaling. PAK proteins serve as targets for the small GTP binding proteins Cdc42 and RAC and have been implicated in a wide range of biological activities. PAK3 forms an activated complex with GTP-bound RAS-like (P21), CDC2 and RAC1 proteins which then catalyzes a variety of targets. Serine/threonine protein kinase that plays a role in a variety of different signaling pathways including cytoskeleton regulation, cell migration, or cell cycle regulation. Plays a role in dendrite spine morphogenesis as well as synapse formation and plasticity. Acts as downstream effector of the small GTPases CDC42 and RAC1. Activation by the binding of active CDC42 and RAC1 results in a conformational change and a subsequent autophosphorylation on several serine and/or threonine residues. Phosphorylates MAPK4 and MAPK6 and activates the downstream target MAPKAPK5, a regulator of F-actin polymerization and cell migration. Additionally, phosphorylates TNNI3/troponin I to modulate calcium sensitivity and relaxation kinetics of thin myofilaments. May also be involved in early neuronal development. In hippocampal neurons, necessary for the formation of dendritic spines and excitatory synapses; this function is dependent on kinase activity and may be exerted by the regulation of actomyosin contractility through the phosphorylation of myosin II regulatory light chain (MLC) (By similarity).

研究领域 (Research Area)

Neuroscience

图片 (Image Data)



Western blot analysis of PAK3 expression in human fetal brain lysate.

注意事项 (Note)

For research use only .