产品名称: Rho A (11B6) Rabbit Monoclonal Antibody

产品货号: AMRe17118



## 产品概述 (Summary)

产品名称 (Production Name) Rho A (11B6) Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit

**应用 (Application)** WB,ICC/IF,FC

种属反应性 (Reactivity) Human, Mouse, Rat

## 产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% New

储存溶液 (Buffer) type preservative N and 50% glycerol. Store at +4°C short term. Store at -

20°C long term. Avoid freeze / thaw cycle.

纯化方式 (Purification) Affinity purification

## 免疫原信息 (Immunogen)

基因名 (Gene Name) RHOA

别名 (Alternative Names) RHOA; ARH12; ARHA; RHO12; RHOH12;

**基因 ID (Gene ID)** 387.0 **蛋白 ID (SwissProt ID)** P61586.

# 产品应用(Application)

**稀释比 (Dilution Ratio)** WB 1:1000-1:5000,ICC/IF 1:100-1:200,FC 1:100-1:200

蛋白分子量 (Molecular Weight) 22kDa

## 研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: Rho A (11B6) Rabbit Monoclonal Antibody

产品货号: AMRe17118



Rho A is a small G protein of the Rho family. Regulates a signal transduction pathway linking plasma membrane receptors to the assembly of focal adhesions and actin stress fibers. Small GTPase which cycles between an active GTP-bound and an inactive GDP-bound state. Mainly associated with cytoskeleton organization, in active state binds to a variety of effector proteins to regulate cellular responses such as cytoskeletal dynamics, cell migration and cell cycle. Regulates a signal transduction pathway linking plasma membrane receptors to the assembly of focal adhesions and actin stress fibers (PubMed:<a href="http://www.uniprot.org/citations/8910519" target=" blank">8910519</a>, PubMed:<a href="http://www.uniprot.org/citations/9121475" target=" blank">9121475</a>, PubMed:<a href="http://www.uniprot.org/citations/31570889" target=" blank">31570889</a>). Involved in a microtubule-dependent signal that is required for the myosin contractile ring formation during cell cycle cytokinesis (PubMed: <a href="mailto:kg."><a href="mailto:kg.">kg.</a> cytokinesis (PubMed: <a href="mailto:kg.") cytokinesis (PubMed: <a href="mailto:kg.">kg.</a href="http://www.uniprot.org/citations/16236794" target=" blank">16236794</a>, PubMed: <a href="http://www.uniprot.org/citations/12900402" target=" blank">12900402</a>). Plays an essential role in cleavage furrow formation. Required for the apical junction formation of keratinocyte cell-cell adhesion (PubMed: <a href="http://www.uniprot.org/citations/20974804" target=" blank">20974804</a>, PubMed:<a href="http://www.uniprot.org/citations/23940119" target=" blank">23940119</a>). Essential for the SPATA13-mediated regulation of cell migration and adhesion assembly and disassembly (PubMed: <a href="http://www.uniprot.org/citations/19934221" target=" blank">19934221</a>). The MEMO1-RHOA-DIAPH1 signaling pathway plays an important role in ERBB2- dependent stabilization of microtubules at the cell cortex. It controls the localization of APC and CLASP2 to the cell membrane, via the regulation of GSK3B activity. In turn, membrane-bound APC allows the localization of the MACF1 to the cell membrane, which is required for microtubule capture and stabilization (PubMed: <a href="http://www.uniprot.org/citations/20937854" target=" blank" > 20937854 </a>). Regulates KCNA2 potassium channel activity by reducing its location at the cell surface in response to CHRM1 activation; promotes KCNA2 endocytosis (PubMed: <a href="http://www.uniprot.org/citations/9635436" target=" blank" > 9635436 </a>, PubMed: <a href="http://www.uniprot.org/citations/19403695" target=" blank">19403695</a>). Acts as an allosteric activator of quanine nucleotide exchange factor ECT2 by binding in its activated GTP-bound form to the PH domain of ECT2 which stimulates the release of PH inhibition and promotes the binding of substrate RHOA to the ECT2 catalytic center (PubMed: <a href="http://www.uniprot.org/citations/31888991" target=" blank">31888991</a>). May be an activator of PLCE1 (PubMed:<a href="http://www.uniprot.org/citations/16103226" target=" blank">16103226</a>). In neurons, involved in the inhibiton of the initial spine growth. Upon activation by CaMKII, modulates dendritic spine structural plasticity by relaying CaMKII transient activation to synapse-specific, long-term signaling (By similarity). Acts as a regulator of platelet alpha-granule release during activation and aggregation of platelets (By similarity).

#### 研究领域 (Research Area)

**Signal Transduction** 


#### 图片 (Image Data)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: Rho A (11B6) Rabbit Monoclonal Antibody

产品货号: AMRe17118





Western blot analysis of Rho A expression in (1) HeLa cell lysate; (2) NIH/3T3 cell lysate.

# 注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838