

产品货号: AMRe19909



产品概述 (Summary)

产品名称 (Production Name) WNK1 (11F15) Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host)Rabbit应用 (Application)WB,ICC/IF种属反应性 (Reactivity)Human

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% New

储存溶液 (Buffer) type preservative N and 50% glycerol. Store at +4°C short term. Store at -

20°C long term. Avoid freeze / thaw cycle.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) WNK1

别名 (Alternative Names) KDP; PSK; p65; HSN2; HSAN2; PRKWNK1;

基因 ID (Gene ID) 65125.0 **蛋白 ID (SwissProt ID)** Q9H4A3.

产品应用(Application)

稀释比 (Dilution Ratio) WB 1:1000-1:5000,ICC/IF 1:50-1:100

蛋白分子量 (Molecular Weight) 251kDa

研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: WNK1 (11F15) Rabbit Monoclonal Antibody


产品货号: AMRe19909



Serine/threonine kinase which plays an important role in the regulation of electrolyte homeostasis, cell signaling, survival, and proliferation. Acts as an activator and inhibitor of sodium-coupled chloride cotransporters and potassium-coupled chloride cotransporters respectively. Activates SCNN1A, SCNN1B, SCNN1D and SGK1. Serine/threonine kinase which plays an important role in the regulation of electrolyte homeostasis, cell signaling, survival, and proliferation. Acts as an activator and inhibitor of sodium-coupled chloride cotransporters and potassium-coupled chloride cotransporters respectively. Activates SCNN1A, SCNN1B, SCNN1D and SGK1. Controls sodium and chloride ion transport by inhibiting the activity of WNK4, by either phosphorylating the kinase or via an interaction between WNK4 and the autoinhibitory domain of WNK1. WNK4 regulates the activity of the thiazide-sensitive Na-Cl cotransporter, SLC12A3, by phosphorylation. WNK1 may also play a role in actin cytoskeletal reorganization. Phosphorylates NEDD4L. Acts as a scaffold to inhibit SLC4A4, SLC26A6 as well as CFTR activities and surface expression, recruits STK39 which mediates the inhibition (By similarity).

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of WNK1 expression in Saos-2 cell lysate.

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838