

产品名称: STAT2 Rabbit Monoclonal Antibody
产品货号: AMRe21125



产品概述 (Summary)

产品名称 (Production Name)	STAT2 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG,Kappa
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein
纯化方式 (Purification)	Protein A

免疫原信息 (Immunogen)

基因名 (Gene Name)	STAT2
别名 (Alternative Names)	STAT2;Signal transducer and activator of transcription 2;p113
基因 ID (Gene ID)	6773.0
蛋白 ID (SwissProt ID)	P52630.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:2000-1:10000,IHC 1:200-1:1000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
蛋白分子量 (Molecular Weight)	Calculated MW:98kD;Observed MW:113kD

研究背景 (Background)

Cell localization:Cytoplasm, Nucleus.The protein encoded by this gene is a member of the STAT protein family. In response


产品名称: STAT2 Rabbit Monoclonal Antibody
产品货号: AMRe21125



to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. In response to interferon (IFN), this protein forms a complex with STAT1 and IFN regulatory factor family protein p48 (ISGF3G), in which this protein acts as a transactivator, but lacks the ability to bind DNA directly. Transcription adaptor P300/CBP (EP300/CREBBP) has been shown to interact specifically with this protein, which is thought to be involved in the process of blocking IFN-alpha response by adenovirus. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Mar 2010],

研究领域 (Research Area)

图片 (Image Data)



Hela cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with primary antibody 1:1000. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

注意事项 (Note)

For research use only .