

产品名称: STAT5b Rabbit Monoclonal Antibody
产品货号: AMRe21131



产品概述 (Summary)

产品名称 (Production Name)	STAT5b Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG,Kappa
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein
纯化方式 (Purification)	Protein A

免疫原信息 (Immunogen)

基因名 (Gene Name)	STAT5B
别名 (Alternative Names)	Signal transducer and activator of transcription 5B
基因 ID (Gene ID)	6777.0
蛋白 ID (SwissProt ID)	P51692.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:2000-1:10000,IHC 1:1000-1:5000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
蛋白分子量 (Molecular Weight)	Calculated MW:90kD;Observed MW:90kD

研究背景 (Background)

Cell localization:Cytoplasm, Nucleus.The protein encoded by this gene is a member of the STAT family of transcription


产品名称: STAT5b Rabbit Monoclonal Antibody
产品货号: AMRe21131



factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein mediates the signal transduction triggered by various cell ligands, such as IL2, IL4, CSF1, and different growth hormones. It has been shown to be involved in diverse biological processes, such as TCR signaling, apoptosis, adult mammary gland development, and sexual dimorphism of liver gene expression. This gene was found to fuse to retinoic acid receptor-alpha (RARA) gene in a small subset of acute promyelocytic leukemias (APLL). The dysregulation of the signaling pathways mediated by this protein may be the cause of the APLL. [provi

研究领域 (Research Area)

图片 (Image Data)



A549 cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with primary antibody 1:1000. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

注意事项 (Note)

For research use only .