

产品名称: AIFM1 Rabbit Monoclonal Antibody
产品货号: AMRe21174



产品概述 (Summary)

产品名称 (Production Name)	AIFM1 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG,Kappa
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein
纯化方式 (Purification)	Protein A

免疫原信息 (Immunogen)

基因名 (Gene Name)	AIFM1
别名 (Alternative Names)	AIFM1;AIF;PDCD8;Apoptosis-inducing factor 1;mitochondrial;Programmed cell death protein 8
基因 ID (Gene ID)	9131.0
蛋白 ID (SwissProt ID)	O95831.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:2000-1:10000,IHC 1:200-1:1000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
蛋白分子量 (Molecular Weight)	Calculated MW:67kD;Observed MW:67kD


研究背景 (Background)

产品名称: AIFM1 Rabbit Monoclonal Antibody
产品货号: AMRe21174

Cell localization: Mitochondrion intermembrane space . Mitochondrion inner membrane. Cytoplasm . Nucleus . Cytoplasm, perinuclear region . Proteolytic cleavage during or just after translocation into the mitochondrial intermembrane space (IMS) results in the formation of an inner-membrane-anchored mature form (AIFmit). During apoptosis, further proteolytic processing leads to a mature form, which is confined to the mitochondrial IMS in a soluble form (AIFsol). AIFsol is released to the cytoplasm in response to specific death signals, and translocated to the nucleus, where it induces nuclear apoptosis (PubMed:15775970). Colocalizes with EIF3G in the nucleus and perinuclear region (PubMed:17094969). .; [Isoform 3]: Mitochondrion intermembrane space . Mitochondrion inner membrane . Has a stronger membrane anchorage than isoform 1. .; [Isoform 4]: Mitochondrion . Cytoplasm, cytosol . In pro-apoptotic conditions, is released from mitochondria to cytosol in a calpain/cathepsin-dependent manner. .; [Isoform 5]: Cytoplasm ..This gene encodes a flavoprotein essential for nuclear disassembly in apoptotic cells, and it is found in the mitochondrial intermembrane space in healthy cells. Induction of apoptosis results in the translocation of this protein to the nucleus where it affects chromosome condensation and fragmentation. In addition, this gene product induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. Mutations in this gene cause combined oxidative phosphorylation deficiency 6 (COXPD6), a severe mitochondrial encephalomyopathy, as well as Cowchock syndrome, also known as X-linked recessive Charcot-Marie-Tooth disease-4 (CMTX-4), a disorder resulting in neuropathy, and axonal and motor-sensory defects with deafness and mental retardation. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome

研究领域 (Research Area)

图片 (Image Data)



Jurkat whole cell lysates were separated by 10% SDS-PAGE, and the membrane was blotted with primary antibody(1:1000). The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

注意事项 (Note)

For research use only .