

产品名称: MEK4 Rabbit Monoclonal Antibody
产品货号: AMRe21291



产品概述 (Summary)

产品名称 (Production Name)	MEK4 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ICC/IF,ELISA,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG,Kappa
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein
纯化方式 (Purification)	Protein A

免疫原信息 (Immunogen)

基因名 (Gene Name)	MAP2K4
别名 (Alternative Names)	MAP2K4;JNKK1;MEK4;MKK4;PRKMK4;SEK1;SERK1;SKK1;Dual specificity mitogen-activated protein kinase kinase 4;MAP kinase kinase 4;MAPKK 4;JNK-activating kinase 1;MAPK/ERK kinase 4;MEK 4;SAPK/ERK kinase 1;SEK1;Stress-activated pro
基因 ID (Gene ID)	6416.0
蛋白 ID (SwissProt ID)	P45985.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:2000-1:10000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
蛋白分子量 (Molecular Weight)	Calculated MW:44kD;Observed MW:44kD

产品名称: MEK4 Rabbit Monoclonal Antibody
产品货号: AMRe21291




研究背景 (Background)

Cell localization: Cytoplasm, Nucleus. This gene encodes a member of the mitogen-activated protein kinase (MAPK) family. Members of this family act as an integration point for multiple biochemical signals and are involved in a wide variety of cellular processes such as proliferation, differentiation, transcription regulation, and development. They form a three-tiered signaling module composed of MAPKKKs, MAPKKs, and MAPKs. This protein is phosphorylated at serine and threonine residues by MAPKKKs and subsequently phosphorylates downstream MAPK targets at threonine and tyrosine residues. A similar protein in mouse has been reported to play a role in liver organogenesis. A pseudogene of this gene is located on the long arm of chromosome X. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2013],

研究领域 (Research Area)

图片 (Image Data)



A431 cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with primary antibody 1:1000. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

注意事项 (Note)

For research use only .