产品名称: MEK3 Rabbit Monoclonal Antibody

产品货号: AMRe21303



产品概述 (Summary)

产品名称 (Production Name) MEK3 Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA,IP

种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

個联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified 同种型 (Isotype) IgG,Kappa 内隆 (Clonality) Monoclonal Liquid

形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein

纯化方式 (Purification) Protein A

免疫原信息 (Immunogen)

基因名 (Gene Name) MAP2K3

MAP2K3;MEK3;MKK3;PRKMK3;SKK2;Dual specificity mitogen-activated

protein kinase kinase 3;MAP kinase kinase 3;MAPKK 3;MAPK/ERK kinase

别名 (Alternative Names)3;MEK 3;Stress-activated protein kinase kinase 2;SAPK kinase 2;SAPKK-

2;SAPKK2

基因 ID (Gene ID) 5606 蛋白 ID (SwissProt ID) P46734.

产品应用 (Application)

WB 1:2000-1:10000,IHC 1:200-1:1000,ICC/IF 1:200-1:1000,ELISA 1:5000-稀释比 (Dilution Ratio)

1:20000,IP 1:50-1:200

蛋白分子量 (Molecular Weight) Calculated MW:39kD;Observed MW:39kD

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: MEK3 Rabbit Monoclonal Antibody

产品货号: AMRe21303




研究背景 (Background)

Cell localization:Cytoplasm, Nucleus.The protein encoded by this gene is a dual specificity protein kinase that belongs to the MAP kinase kinase family. This kinase is activated by mitogenic and environmental stress, and participates in the MAP kinase-mediated signaling cascade. It phosphorylates and thus activates MAPK14/p38-MAPK. This kinase can be activated by insulin, and is necessary for the expression of glucose transporter. Expression of RAS oncogene is found to result in the accumulation of the active form of this kinase, which thus leads to the constitutive activation of MAPK14, and confers oncogenic transformation of primary cells. The inhibition of this kinase is involved in the pathogenesis of Yersina pseudotuberculosis. Multiple alternatively spliced transcript variants that encode distinct isoforms have been reported for this gene. [provided by RefSeq, Jul 2008],

研究领域 (Research Area)

图片 (Image Data)



Jurkat cell lysates were separated by 4-20% SDS-PAGE, and the membrane was blotted with primary antibody 1:1000. The HRP-conjugated Goat anti-Rabbit IgG(H + L) antibody was used to detect the antibody.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838