

产品名称: NQO1 Rabbit Monoclonal Antibody
产品货号: AMRe21489



产品概述 (Summary)

产品名称 (Production Name)	NQO1 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG,Kappa
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%protective protein
纯化方式 (Purification)	Protein A

免疫原信息 (Immunogen)

基因名 (Gene Name)	NQO1
别名 (Alternative Names)	NQO1;DIA4;NMOR1;NAD;P;H dehydrogenase [quinone] 1;Azoreductase;DT-diaphorase;DTD;Menadione reductase;NAD;P;H:quinone oxidoreductase 1;Phylloquinone reductase;Quinone reductase 1;QR1
基因 ID (Gene ID)	1728.0
蛋白 ID (SwissProt ID)	P15559.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000,IHC 1:200-1:1000,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
蛋白分子量 (Molecular Weight)	Calculated MW:31kD;Observed MW:31kD

产品名称: NQO1 Rabbit Monoclonal Antibody
产品货号: AMRe21489




研究背景 (Background)


Cell localization: Cytoplasm. This gene is a member of the NAD(P)H dehydrogenase (quinone) family and encodes a cytoplasmic 2-electron reductase. This FAD-binding protein forms homodimers and reduces quinones to hydroquinones. This protein's enzymatic activity prevents the one electron reduction of quinones that results in the production of radical species. Mutations in this gene have been associated with tardive dyskinesia (TD), an increased risk of hematotoxicity after exposure to benzene, and susceptibility to various forms of cancer. Altered expression of this protein has been seen in many tumors and is also associated with Alzheimer's disease (AD). Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008],

研究领域 (Research Area)

图片 (Image Data)




Western blot analysis of lysates from Hela cells, using NQO1 Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.




Immunohistochemical analysis of paraffin-embedded Human kidney tissue. 1, primary Antibody was diluted at 1:200(4°C, overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C, 20min). 3, Secondary antibody was diluted at 1:200(room temperature, 30min).

产品名称: NQO1 Rabbit Monoclonal Antibody
产品货号: AMRe21489



Immunohistochemical analysis of paraffin-embedded Mouse kidney tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Rat kidney tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

注意事项 (Note)

For research use only .