

产品名称: METTL3 Rabbit Monoclonal Antibody
产品货号: AMRe21565



产品概述 (Summary)

产品名称 (Production Name)	METTL3 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG,Kappa
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein
纯化方式 (Purification)	Protein A

免疫原信息 (Immunogen)

基因名 (Gene Name)	METTL3
别名 (Alternative Names)	N6-adenosine-methyltransferase 70 kDa subunit;MT-A70;Methyltransferase-like protein 3;
基因 ID (Gene ID)	56339.0
蛋白 ID (SwissProt ID)	Q86U44.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000,IHC 1:200-1:400,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000,IP 1:50-1:200
蛋白分子量 (Molecular Weight)	Calculated MW:64kD;Observed MW:70kD

研究背景 (Background)


产品名称: METTL3 Rabbit Monoclonal Antibody
产品货号: AMRe21565



Cell localization:Nucleus.This gene encodes the 70 kDa subunit of MT-A which is part of N6-adenosine-methyltransferase. This enzyme is involved in the posttranscriptional methylation of internal adenosine residues in eukaryotic mRNAs, forming N6-methyladenosine. [provided by RefSeq, Jul 2008]


研究领域 (Research Area)

图片 (Image Data)




Western blot analysis of lysates from NCCIT

cells, using METTL3 Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.



Immunohistochemical analysis of paraffin-embedded mouse testis tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

产品名称: METTL3 Rabbit Monoclonal Antibody
产品货号: AMRe21565



Immunohistochemical analysis of paraffin-embedded prostate cancer tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

注意事项 (Note)

For research use only .