产品名称: Lactate Dehydrogenase Isoenzyme V Rabbit

Monoclonal Antibody 产品货号: AMRe21598



产品概述 (Summary)

产品名称 (Production Name) Lactate Dehydrogenase Isoenzyme V Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,ELISA,IP种属反应性 (Reactivity)Human,Mouse,Rat

产品性能 (Performance)

(はいけん) は Unconjugated (Modification) Unmodified (Isotype) は IgG,Kappa 内隆 (Clonality) Monoclonal

形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

储存溶液 (Buffer) PBS, 50% glycerol, 0.05% Proclin 300, 0.05% protective protein

纯化方式 (Purification) Protein A

免疫原信息 (Immunogen)

基因名 (Gene Name) LDHA PIG19

别名 (Alternative Names)

基因 ID (Gene ID) 3939.0 **蛋白 ID (SwissProt ID)** P00338.

产品应用(Application)

WB 1:500-1:2000,IHC 1:200-1:1000,ICC/IF 1:200-1:1000,ELISA 1:5000-**稀释比 (Dilution Ratio)**

1:20000,IP 1:50-1:200

蛋白分子量 (Molecular Weight) Calculated MW:36kD;Observed MW:36kD

研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: Lactate Dehydrogenase Isoenzyme V Rabbit


Monoclonal Antibody 产品货号: AMRe21598



Cell localization:Cytoplasm.The protein encoded by this gene catalyzes the conversion of L-lactate and NAD to pyruvate and NADH in the final step of anaerobic glycolysis. The protein is found predominantly in muscle tissue and belongs to the lactate dehydrogenase family. Mutations in this gene have been linked to exertional myoglobinuria. Multiple transcript variants encoding different isoforms have been found for this gene. The human genome contains several non-transcribed pseudogenes of this gene. [provided by RefSeq, Sep 2008],


研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of lysates from A431

cells, using Lactate Dehydrogenase Isoenzyme V Rabbit mAb. The HRP-conjugated Goat anti-Rabbit IgG antibody was used to detect the antibody.



Immunohistochemical analysis of paraffin-embedded human appendix tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

Monoclonal Antibody 产品货号: AMRe21598





Immunohistochemical analysis of paraffin-embedded mouse heart tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded rat heart tissue. 1, primary Antibody was diluted at 1:200(4°C,overnight). 2, EDTA pH 9.0 was used for antibody retrieval(>98°C,20min). 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

注意事项 (Note)

For research use only.

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838