

产品名称: HDAC11 Rabbit Monoclonal Antibody
产品货号: AMRe83684



产品概述 (Summary)

产品名称 (Production Name)	HDAC11 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IP
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide, 0.05% protective protein and 50% glycerol.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	HDAC11
别名 (Alternative Names)	HDAC11; HD11; Histone deacetylase 11;;HDAC11
基因 ID (Gene ID)	
蛋白 ID (SwissProt ID)	Q96DB2.A synthesized peptide derived from human HDAC11

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:2000, IP 1:20-1:50
蛋白分子量 (Molecular Weight)	39 kDa

研究背景 (Background)

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4).


产品名称: HDAC11 Rabbit Monoclonal Antibody
产品货号: AMRe83684




Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes.

研究领域 (Research Area)


图片 (Image Data)



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot analysis of HDAC11 expression in (1) Jurkat cell lysate;(2) MCF7 cell lysate.

产品名称: HDAC11 Rabbit Monoclonal Antibody
产品货号: AMRe83684



注意事项 (Note)

For research use only .