

产品名称: WDR4 Rabbit Monoclonal Antibody
产品货号: AMRe84032



产品概述 (Summary)

产品名称 (Production Name)	WDR4 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ICC,FC
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in PBS with 0.05% sodium azide,0.05% protective protein and 50% glycerol.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	WDR4
别名 (Alternative Names)	TRM82; WD repeat protein 4; wdr4;;WDR4
基因 ID (Gene ID)	
蛋白 ID (SwissProt ID)	P57081.A synthesized peptide derived from human WDR4

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:2000,IHC 1:100-1:200,ICC/IF 1:50-1:200,ICC 1:50-1:200,FC 1:20-1:100
蛋白分子量 (Molecular Weight)	45 kDa

研究背景 (Background)


产品名称: WDR4 Rabbit Monoclonal Antibody
产品货号: AMRe84032




Required for the formation of N(7)-methylguanine at position 46 (m7G46) in tRNA. In the complex, it is required to stabilize and induce conformational change of the catalytic subunit.

研究领域 (Research Area)


图片 (Image Data)



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.

产品名称: WDR4 Rabbit Monoclonal Antibody
产品货号: AMRe84032



Immunofluorescent analysis using the Antibody at 1:50 dilution.

注意事项 (Note)

For research use only .