

产品名称: Cullin 4A/4B Rabbit Monoclonal Antibody
产品货号: AMRe85466



产品概述 (Summary)

产品名称 (Production Name)	Cullin 4A/4B Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC
种属反应性 (Reactivity)	Human,Rat,Hamster

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in TBS with 0.05% sodium azide,0.05%protective protein and 50% glycerol.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	Cullin 4A/4B
别名 (Alternative Names)	CUL4B; KIAA0695; Cullin-4B; CUL-4B
基因 ID (Gene ID)	8450.0
蛋白 ID (SwissProt ID)	Q13620.A synthetic peptide of human Cullin 4B

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000,IHC 1:50-1:100
蛋白分子量 (Molecular Weight)	Calculated MW: 104 kDa; Observed MW: 88,104 kDa

研究背景 (Background)

Core component of multiple cullin-RING-based E3 ubiquitin-protein ligase complexes which mediate the ubiquitination


产品名称: Cullin 4A/4B Rabbit Monoclonal Antibody
产品货号: AMRe85466




and subsequent proteasomal degradation of target proteins. The functional specificity of the E3 ubiquitin-protein ligase complex depends on the variable substrate recognition subunit. CUL4B may act within the complex as a scaffold protein, contributing to catalysis through positioning of the substrate and the ubiquitin-conjugating enzyme. Plays a role as part of the E3 ubiquitin-protein ligase complex in polyubiquitination of CDT1, histone H2A, histone H3 and histone H4 in response to radiation-induced DNA damage. Targeted to UV damaged chromatin by DDB2 and may be important for DNA repair and DNA replication. Required for ubiquitination of cyclin E, and consequently, normal G1 cell cycle progression. Regulates the mammalian target-of-rapamycin (mTOR) pathway involved in control of cell growth, size and metabolism. Specific CUL4B regulation of the mTORC1-mediated pathway is dependent upon 26S proteasome function and requires interaction between CUL4B and MLST8.

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of Cullin 4A/CUL4A+Cullin 4B/CUL4B in Jurkat, C6, CHO-K1 lysates using Cullin 4A/4B antibody.



Immunohistochemistry analysis of paraffin-embedded Human lung cancer using Cullin 4A/B antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.

产品名称: Cullin 4A/4B Rabbit Monoclonal Antibody
产品货号: AMRe85466



注意事项 (Note)

For research use only .