

产品名称: DBC 1 Rabbit Monoclonal Antibody
产品货号: AMRe85492



产品概述 (Summary)

产品名称 (Production Name)	DBC 1 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC,IP
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in TBS with 0.05% sodium azide, 0.05% protective protein and 50% glycerol.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	DBC 1
别名 (Alternative Names)	DBC1; DBC-1; NET35; p30DBC; p30 DBC; KIAA1967
基因 ID (Gene ID)	57805.0
蛋白 ID (SwissProt ID)	Q8N163.A synthetic peptide of human DBC-1

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000, IHC 1:50-1:100, ICC 1:50-1:200, IP 1:10-1:20
蛋白分子量 (Molecular Weight)	Calculated MW: 103 kDa; Observed MW: 130 kDa

研究背景 (Background)

Core component of the DBIRD complex, a multiprotein complex that acts at the interface between core mRNP particles and

产品名称: DBC 1 Rabbit Monoclonal Antibody
产品货号: AMRe85492




RNA polymerase II (RNAPII) and integrates transcript elongation with the regulation of alternative splicing; the DBIRD complex affects local transcript elongation rates and alternative splicing of a large set of exons embedded in (A + T)-rich DNA regions. Inhibits SIRT1 deacetylase activity leading to increasing levels of p53/TP53 acetylation and p53-mediated apoptosis. Inhibits SUV39H1 methyltransferase activity. As part of a histone H3-specific methyltransferase complex may mediate ligand-dependent transcriptional activation by nuclear hormone receptors. Plays a critical role in maintaining genomic stability and cellular integrity following UV-induced genotoxic stress. Regulates the circadian expression of the core clock components NR1D1 and ARNTL/BMAL1. Enhances the transcriptional repressor activity of NR1D1 through stabilization of NR1D1 protein levels by preventing its ubiquitination and subsequent degradation (PubMed:18235501, PubMed:18235502, PubMed:19131338, PubMed:19218236, PubMed:22446626, PubMed:23352644, PubMed:23398316). Represses the ligand-dependent transcriptional activation function of ESR2 (PubMed:20074560). Acts as a regulator of PCK1 expression and gluconeogenesis by a mechanism that involves, at least in part, both NR1D1 and SIRT1 (PubMed:24415752). Negatively regulates the deacetylase activity of HDAC3 and can alter its subcellular localization (PubMed:21030595). Positively regulates the beta-catenin pathway (canonical Wnt signaling pathway) and is required for MCC-mediated repression of the beta-catenin pathway (PubMed:24824780). Represses ligand-dependent transcriptional activation function of NR1H2 and NR1H3 and inhibits the interaction of SIRT1 with NR1H3 (PubMed:25661920). Plays an important role in tumor suppression through p53/TP53 regulation; stabilizes p53/TP53 by affecting its interaction with ubiquitin ligase MDM2 (PubMed:25732823). Represses the transcriptional activator activity of BRCA1 (PubMed:20160719). Inhibits SIRT1 in a CHEK2 and PSEM3-dependent manner and inhibits the activity of CHEK2 in vitro (PubMed:25361978).

研究领域 (Research Area)

Apoptosis, Wnt signaling pathway

图片 (Image Data)



Western blot analysis of DBC1 in 293, Jurkat lysates using DBC 1 antibody.

注意事项 (Note)

For research use only .