

产品名称: KAT2A Rabbit Monoclonal Antibody
产品货号: AMRe85602



产品概述 (Summary)

产品名称 (Production Name)	KAT2A Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IP
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Purified antibody in TBS with 0.05% sodium azide, 0.05% protective protein and 50% glycerol.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	KAT2A
别名 (Alternative Names)	GCN5; hGCN5; GCN5L2; PCAF-b
基因 ID (Gene ID)	2648.0
蛋白 ID (SwissProt ID)	Q92830.Recombinant protein of human GCN5

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:1000, IP 1:10-1:20
蛋白分子量 (Molecular Weight)	Calculated MW: 94 kDa; Observed MW: 94 kDa

研究背景 (Background)

Protein lysine acyltransferase that can act both as a acetyltransferase and succinyltransferase, depending on the context


产品名称: KAT2A Rabbit Monoclonal Antibody
产品货号: AMRe85602



(PubMed:29211711). Acts as a histone lysine succinyltransferase: catalyzes succinylation of histone H3 on 'Lys-79' (H3K79succ), with a maximum frequency around the transcription start sites of genes (PubMed:29211711). Succinylation of histones gives a specific tag for epigenetic transcription activation (PubMed:29211711). Association with the 2-oxoglutarate dehydrogenase complex, which provides succinyl-CoA, is required for histone succinylation (PubMed:29211711). In different complexes, functions either as an acetyltransferase (HAT) or as a succinyltransferase: in the SAGA and ATAC complexes, acts as a histone acetyltransferase (PubMed:17301242, PubMed:19103755, PubMed:29211711). Has significant histone acetyltransferase activity with core histones, but not with nucleosome core particles (PubMed:17301242, PubMed:19103755). Acetylation of histones gives a specific tag for epigenetic transcription activation (PubMed:17301242, PubMed:19103755, PubMed:29211711). Involved in long-term memory consolidation and synaptic plasticity: acts by promoting expression of a hippocampal gene expression network linked to neuroactive receptor signaling . Acts as a positive regulator of T-cell activation: upon TCR stimulation, recruited to the IL2 promoter following interaction with NFATC2 and catalyzes acetylation of histone H3 at Lys-9 (H3K9ac), leading to promote IL2 expression . Also acetylates non-histone proteins, such as CEBPB, PLK4 and TBX5 (PubMed:17301242, PubMed:29174768, PubMed:27796307). Involved in heart and limb development by mediating acetylation of TBX5, acetylation regulating nucleocytoplasmic shuttling of TBX5 (PubMed:29174768). Acts as a negative regulator of centrosome amplification by mediating acetylation of PLK4 (PubMed:27796307).

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of KAT2A / GCN5 in K562 lysates using KAT2A antibody.

注意事项 (Note)

For research use only .