产品名称: LIM Kinase 1 Rabbit Monoclonal Antibody

产品货号: AMRe86557



产品概述 (Summary)

产品名称 (**Production Name**) LIM Kinase 1 Rabbit Monoclonal Antibody

描述 (**Description**) Recombinant rabbit monoclonal antibody

宿主 (Host)Rabbit应用 (Application)WB种属反应性 (Reactivity)Human

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01%

储存溶液 (Buffer) sodium azide and 0.05% protective protein. Stable for 12 months from date

of receipt.

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name) LIM Kinase 1 别名 (Alternative Names) LIMK; LIMK-1

基因 ID (Gene ID) 3984 **蛋白 ID (SwissProt ID)** P53667.

产品应用(Application)

稀释比 (Dilution Ratio) WB 1:1000-1:5000

蛋白分子量 (Molecular Weight) Calculated MW:73 kDa; Observed MW:73 kDa

研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838




产品货号: AMRe86557



There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. LIMK1 is a serine/threonine kinase that regulates actin polymerization via phosphorylation and inactivation of the actin binding factor cofilin. This protein is ubiquitously expressed during development and plays a role in many cellular processes associated with cytoskeletal structure. This protein also stimulates axon growth and may play a role in brain development. LIMK1 hemizygosity is implicated in the impaired visuospatial constructive cognition of Williams syndrome. Alternative splicing results in multiple transcript variants encoding distinct isoforms.[provided by RefSeq, Feb 2011]

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from U-87MG cells using AMRe86557 at 1:1000.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838