产品名称: p21 Rabbit Monoclonal Antibody

产品货号: AMRe86606



产品概述 (Summary)

产品名称 (Production Name) p21 Rabbit Monoclonal Antibody

描述 (**Description**) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,FC,IP

种属反应性 (Reactivity) Human

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01%

储存溶液 (Buffer) sodium azide and 0.05% protective protein. Stable for 12 months from date

of receipt.

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name) p21

别名 (Alternative Names) P21; CIP1; SDI1; WAF1; CAP20; CDKN1; MDA-6; p21CIP1

基因 ID (Gene ID) 1026 **蛋白 ID (SwissProt ID)** P38936.

产品应用(Application)

WB 1:1000-1:5000,IHC 1:100-1:200,ICC/IF 1:500-1:1000,FC 1:50-1:200,IP 1:10-

稀释比 (Dilution Ratio)
1:100

蛋白分子量 (Molecular Weight) Calculated MW:18 kDa; Observed MW:21 kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: p21 Rabbit Monoclonal Antibody

产品货号: AMRe86606




研究背景 (Background)

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or -cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at G1. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Multiple alternatively spliced variants have been found for this gene. [provided by RefSeq, Sep 2015]

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from HeLa, HCT116 cells using AMRe86606 at 1:1000.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838