产品名称: TRAF2 Rabbit Monoclonal Antibody

产品货号: AMRe86844



产品概述 (Summary)

产品名称 (Production Name) TRAF2 Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,FC,IP种属反应性 (Reactivity)Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01%

储存溶液 (Buffer) sodium azide and 0.05% protective protein. Stable for 12 months from date

of receipt.

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name) TRAF2

別名 (Alternative Names) TRAP; TRAP3; MGC:45012

基因 ID (Gene ID) 7186 **蛋白 ID (SwissProt ID)** Q12933.

产品应用(Application)

WB 1:500-1:2000,IHC 1:100-1:200,ICC/IF 1:100-1:200,FC 1:50-1:200,IP 1:20-

稀释比 (Dilution Ratio) 1:50

蛋白分子量 (Molecular Weight) Calculated MW:56 kDa; Observed MW:53 kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: TRAF2 Rabbit Monoclonal Antibody

产品货号: AMRe86844




研究背景 (Background)


The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alphamediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can unbiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis. Multiple alternatively spliced transcript variants have been found for this gene, but the biological validity of only one transcript has been determined. [provided by RefSeq, Jul 2008]

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from THP-1, HEK293T, HeLa cells using AMRe86844 at 1:1000.



Western blot analysis of extracts from MOLT-4 cells using AMRe86844 at 1:1000.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: TRAF2 Rabbit Monoclonal Antibody

产品货号: AMRe86844



注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838