产品名称: PAX5 Rabbit Monoclonal Antibody

产品货号: AMRe86891



## 产品概述 (Summary)

产品名称 (Production Name) PAX5 Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,FC,IP种属反应性 (Reactivity)Human,Mouse,Rat

### 产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01%

**储存溶液 (Buffer)** sodium azide and 0.05% protective protein. Stable for 12 months from date

of receipt.

纯化方式 (Purification) Affinity Purification

## 免疫原信息 (Immunogen)

基因名 (Gene Name) PAX5

别名 (Alternative Names) ALL3; BSAP

**基因 ID (Gene ID)** 5079 **蛋白 ID (SwissProt ID)** Q02548.

# 产品应用(Application)

WB 1:1000-1:5000,IHC 1:1000-1:5000,ICC/IF 1:100-1:200,FC 1:20-1:50,IP 1:20-

稀释比 (Dilution Ratio) 1:50

蛋白分子量 (Molecular Weight) Calculated MW:42 kDa; Observed MW:45 kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: PAX5 Rabbit Monoclonal Antibody

产品货号: AMRe86891




#### 研究背景 (Background)

This gene encodes a member of the paired box (PAX) family of transcription factors. The central feature of this gene family is a novel, highly conserved DNA-binding motif, known as the paired box. Paired box transcription factors are important regulators in early development, and alterations in the expression of their genes are thought to contribute to neoplastic transformation. This gene encodes the B-cell lineage specific activator protein that is expressed at early, but not late stages of B-cell differentiation. Its expression has also been detected in developing CNS and testis and so the encoded protein may also play a role in neural development and spermatogenesis. This gene is located at 9p13, which is involved in t(9;14) (p13;q32) translocations recurring in small lymphocytic lymphomas of the plasmacytoid subtype, and in derived large-cell lymphomas. This translocation brings the potent E-mu enhancer of the IgH gene into close proximity of the PAX5 promoter, suggesting that the deregulation of transcription of this gene contributes to the pathogenesis of these lymphomas. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2013]

#### 研究领域 (Research Area)

#### 图片 (Image Data)



Western blot analysis of extracts from Raji cells using AMRe86891 at 1:1000.

#### 注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838