

产品名称: PEG10 Rabbit Monoclonal Antibody
产品货号: AMRe87083



产品概述 (Summary)

产品名称 (Production Name)	PEG10 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF, FC
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% protective protein. Stable for 12 months from date of receipt.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PEG10
别名 (Alternative Names)	EDR; HB-1; Mar2; RTL2; MEF3L; Mart2; RGAG3; SIRH1
基因 ID (Gene ID)	23089
蛋白 ID (SwissProt ID)	Q86TG7.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:1000-1:5000, ICC/IF 1:100-1:200, FC 1:50-1:100
蛋白分子量 (Molecular Weight)	Calculated MW:80 kDa; Observed MW:100,55 kDa

研究背景 (Background)


产品名称: PEG10 Rabbit Monoclonal Antibody
产品货号: AMRe87083



This is a paternally expressed imprinted gene that is thought to have been derived from the Ty3/Gypsy family of retrotransposons. It contains two overlapping open reading frames, RF1 and RF2, and expresses two proteins: a shorter, gag-like protein (with a CCHC-type zinc finger domain) from RF1; and a longer, gag/pol-like fusion protein (with an additional aspartic protease motif) from RF1/RF2 by -1 translational frameshifting (-1 FS). While -1 FS has been observed in RNA viruses and transposons in both prokaryotes and eukaryotes, this gene represents the first example of -1 FS in a eukaryotic cellular gene. This gene is highly conserved across mammalian species and retains the heptanucleotide (GGGAAAC) and pseudoknot elements required for -1 FS. It is expressed in adult and embryonic tissues (most notably in placenta) and reported to have a role in cell proliferation, differentiation and apoptosis. Overexpression of this gene has been associated with several malignancies, such as hepatocellular carcinoma and B-cell lymphocytic leukemia. Knockout mice lacking this gene showed early embryonic lethality with placental defects, indicating the importance of this gene in embryonic development. Additional isoforms resulting from alternatively spliced transcript variants, and use of upstream non-AUG (CUG) start codon have been reported for this gene. [provided by RefSeq, Oct 2014]

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from HepG2 cells using AMRe87083 at 1:1000.

注意事项 (Note)

For research use only .