产品名称: CRF Rabbit Monoclonal Antibody

产品货号: AMRe87333



产品概述 (Summary)

产品名称 (Production Name) CRF Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit 应用 (Application) WB

种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01%

储存溶液 (Buffer) sodium azide and 0.05% protective protein. Stable for 12 months from date

of receipt.

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name) CRF

別名 (Alternative Names)CRF; CRH1基因 ID (Gene ID)1392蛋白 ID (SwissProt ID)P06850.

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000

蛋白分子量 (Molecular Weight) Calculated MW:21 kDa; Observed MW:21 kDa

研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: CRF Rabbit Monoclonal Antibody


产品货号: AMRe87333



This gene encodes a member of the corticotropin-releasing factor family. The encoded preproprotein is proteolytically processed to generate the mature neuropeptide hormone. In response to stress, this hormone is secreted by the paraventricular nucleus (PVN) of the hypothalamus, binds to corticotropin releasing hormone receptors and stimulates the release of adrenocorticotropic hormone from the pituitary gland. Marked reduction in this protein has been observed in association with Alzheimer's disease. Autosomal recessive hypothalamic corticotropin deficiency has multiple and potentially fatal metabolic consequences including hypoglycemia and hepatitis. In addition to production in the hypothalamus, this protein is also synthesized in peripheral tissues, such as T lymphocytes, and is highly expressed in the placenta. In the placenta it is a marker that determines the length of gestation and the timing of parturition and delivery. A rapid increase in circulating levels of the hormone occurs at the onset of parturition, suggesting that, in addition to its metabolic functions, this protein may act as a trigger for parturition. [provided by RefSeq, Nov 2015]

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from Mouse brain tissue using AMRe87333 at 1:1000.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838