

产品名称: BCR Rabbit Monoclonal Antibody
产品货号: AMRe87495



产品概述 (Summary)

产品名称 (Production Name)	BCR Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,FC,IP
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% protective protein. Stable for 12 months from date of receipt.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	BCR
别名 (Alternative Names)	ALL; CML; PHL; BCR1; D22S11; D22S662
基因 ID (Gene ID)	613
蛋白 ID (SwissProt ID)	P11274.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:200-1:500,FC 1:200-1:500,IP 1:20-1:50
蛋白分子量 (Molecular Weight)	Calculated MW:143 kDa; Observed MW:150 kDa

研究背景 (Background)


产品名称: BCR Rabbit Monoclonal Antibody
产品货号: AMRe87495



A reciprocal translocation between chromosomes 22 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from 3T3 cells using AMRe87495 at 1:1000.

注意事项 (Note)

For research use only .