产品名称: EphB2 Rabbit Monoclonal Antibody

产品货号: AMRe87531



产品概述 (Summary)

产品名称 (Production Name) EphB2 Rabbit Monoclonal Antibody

描述 (Description) Recombinant rabbit monoclonal antibody

宿主 (Host) Rabbit 应用 (Application) WB,IHC

种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Monoclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01%

储存溶液 (Buffer) sodium azide and 0.05% protective protein. Stable for 12 months from date

of receipt.

纯化方式 (Purification) Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name) EphB2

别名 (Alternative Names) DRT; EK5; ERK; CAPB; Hek5; PCBC; EPHT3; Tyro5; BDPLT22

基因 ID (Gene ID) 2048 **蛋白 ID (SwissProt ID)** P29323.

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:1000-1:5000,IHC 1:200-1:1000

蛋白分子量 (Molecular Weight) Calculated MW:118 kDa; Observed MW:130 kDa

研究背景 (Background)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: EphB2 Rabbit Monoclonal Antibody


产品货号: AMRe87531



This gene encodes a member of the Eph receptor family of receptor tyrosine kinase transmembrane glycoproteins. These receptors are composed of an N-terminal glycosylated ligand-binding domain, a transmembrane region and an intracellular kinase domain. They bind ligands called ephrins and are involved in diverse cellular processes including motility, division, and differentiation. A distinguishing characteristic of Eph-ephrin signaling is that both receptors and ligands are competent to transduce a signaling cascade, resulting in bidirectional signaling. This protein belongs to a subgroup of the Eph receptors called EphB. Proteins of this subgroup are distinguished from other members of the family by sequence homology and preferential binding affinity for membrane-bound ephrin-B ligands. Allelic variants are associated with prostate and brain cancer susceptibility. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2015]

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from Mouse brain tissue using AMRe87531 at 1:1000.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838