

产品名称: PHD3 Rabbit Monoclonal Antibody
产品货号: AMRe87588



产品概述 (Summary)

产品名称 (Production Name)	PHD3 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF, IP
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% protective protein. Stable for 12 months from date of receipt.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PHD3
别名 (Alternative Names)	PHD3; HIFPH3; HIFP4H3
基因 ID (Gene ID)	112399
蛋白 ID (SwissProt ID)	Q9H6Z9.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000, ICC/IF 1:100-1:200, IP 1:50-1:100
蛋白分子量 (Molecular Weight)	Calculated MW:27 kDa; Observed MW:27 kDa

研究背景 (Background)


产品名称: PHD3 Rabbit Monoclonal Antibody
产品货号: AMRe87588



Cellular oxygen sensor that catalyzes, under normoxic conditions, the post-translational formation of 4-hydroxyproline in hypoxia-inducible factor (HIF) alpha proteins. Hydroxylates a specific proline found in each of the oxygen-dependent degradation (ODD) domains (N-terminal, NODD, and C-terminal, CODD) of HIF1A. Also hydroxylates HIF2A. Has a preference for the CODD site for both HIF1A and HIF2A. Hydroxylation on the NODD site by EGLN3 appears to require prior hydroxylation on the CODD site. Hydroxylated HIFs are then targeted for proteasomal degradation via the von Hippel-Lindau ubiquitination complex. Under hypoxic conditions, the hydroxylation reaction is attenuated allowing HIFs to escape degradation resulting in their translocation to the nucleus, heterodimerization with HIF1B, and increased expression of hypoxia-inducible genes.

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from HeLa cells and Rat spleen tissue using AMRe87588 at 1:1000.

注意事项 (Note)

For research use only .