

产品名称: TGF beta 1/3 Rabbit Monoclonal Antibody
产品货号: AMRe87649



产品概述 (Summary)

产品名称 (Production Name)	TGF beta 1/3 Rabbit Monoclonal Antibody
描述 (Description)	Recombinant rabbit monoclonal antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Monoclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% protective protein. Stable for 12 months from date of receipt.
纯化方式 (Purification)	Affinity Purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TGF beta 1/3
别名 (Alternative Names)	CED; LAP; DPD1; TGFB; TGFbeta
基因 ID (Gene ID)	7040, 7043
蛋白 ID (SwissProt ID)	P01137, P10600.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000
蛋白分子量 (Molecular Weight)	Calculated MW:44,47 kDa; Observed MW:44,12 kDa

研究背景 (Background)


产品名称: TGF beta 1/3 Rabbit Monoclonal Antibody
产品货号: AMRe87649



This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGFB family members. This encoded protein regulates cell proliferation, differentiation and growth, and can modulate expression and activation of other growth factors including interferon gamma and tumor necrosis factor alpha. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease. [provided by RefSeq, Aug 2016]

研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of extracts from K562 cells using AMRe87649 at 1:1000.

注意事项 (Note)

For research use only .