
# Recombinant Human CCL18 (N-6His)

*Catalog#:PEH0262*

## 概述（Summary）

| | |
|---|---|
| **别称（Alternative Names）** | C-C Motif Chemokine 18; Alternative Macrophage Activation-Associated CC Chemokine 1; AMAC-1; CC Chemokine PARC; Dendritic Cell Chemokine 1; DC-CK1; Macrophage Inflammatory Protein 4; MIP-4; Pulmonary and Activation-Regulated Chemokine; Small-Inducible Cytokine A18; CCL18; AMAC1; DCCK1; MIP4; PARC; SCYA18 |
| **纯度（Purity）** | Greater than 95% as determined by reducing SDS-PAGE |
| **内毒素（Endotoxin level）** | <1 EU/μg as determined by LAL test. |
| **蛋白构建（Construction）** | 6His-【Ala21-Ala89】 |
| **Accession #** | P55774 |
| **表达宿主（Host）** | E. coli |
| **种属（Species）** | Human |
| **预测分子量（Predicted MW）** | 10.1 kDa |
| **制剂（Form）** | Lyophilized from a 0.2 μm filtered solution of 20mM Citrate, 6% Trehalose, 4% Mannitol, 0.05% Tween 80, pH 4.0. |
| **运输方式（Shipping）** | The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below. |
| **稳定性&储存（Stability &Storage）** | 12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles. |
| **复溶（Reconstitution）** | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

## 电泳图（SDS-PAGE image）




Greater than 95% as determined by reducing SDS-PAGE.

## 注意事项（Note）

For Research Use Only , Not for Diagnostic Use.