
Recombinant Human IL-17F

Catalog#:PEH0879

概述（Summary）


别称（Alternative Names）	Interleukin-17F; IL-17F; Cytokine ML-1; IL17F
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	【Arg31-Gln163】
Accession #	Q96PD4
表达宿主（Host）	E. coli
种属（Species）	Human
预测分子量（Predicted MW）	15 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM PB, pH 7.4.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

电泳图（SDS-PAGE image）




Greater than 95% as determined by reducing SDS-PAGE.


活性图（Bioactivity image）




Measured by its ability to induce IL-6 secretion by NIH3T3 mouse embryonic fibroblast cells. The ED50 for this effect is 5-40 ng/ml.



Immobilized Human IL-17F at 2μg/ml (100 μl/well)can bind Biotinylated Human IL-17RA-His-Avi.The ED50 of Biotinylated Human IL-17RA-His-Avi is 0.11 ug/ml.



Loaded Biotinylated Human IL-17RA-His-Avi on HIS1K Biosensor, can bind Human IL-17F with an affinity constant of 0.56 uM nM as determined in BLI assay.



Loaded Biotinylated Human IL-17RA-Fc on Pro A Biosensor, can bind Human IL-17F with an affinity constant of 0.15uM as determined in BLI assay.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.