
Active Interleukin 1 Alpha (IL1a)

Catalog#:PEH12110

概述（Summary）

别称（Alternative Names）	IL1-A; IL-1α; IL1F1; Preinterleukin 1 Alpha; Hematopoietin-1; Pro-Interleukin-1-Alpha; Interleukin-1 Family Member 1
纯度（Purity）	Greater than 97% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	Please contact with the lab for this information
蛋白构建（Construction）	【Ser113-Ala271】
Accession #	Q53QF9
表达宿主（Host）	E.coli
种属（Species）	Homo sapiens (Human)
预测分子量（Predicted MW）	21.2 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.

电泳图（SDS-PAGE image）



Greater than 97% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）




Figure 2. Inhibitory effect of IL1α on cell proliferation of MCF-7 cells.

IL1α (Interleukin-1 alpha) is a member of the interleukin 1 cytokine family. This cytokine is produced by monocytes and macrophages as a proprotein, which is proteolytically processed and released in response to cell injury, and thus induces cell apoptosis. It is reported that exposure of MCF-7 cells to certain concentration of IL1α results in inhibition of cell growth. Thus, an cell proliferation assay of MCF-7 was conducted with the addition of IL1α. MCF-7 cells were seeded overnight at a density of 5,000 cells/well, and then treated with or without various concentrations of IL1α for 72h, then cells were observed by inverted microscope and cell viability was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10μL of CCK-8 solution was added to each well of the plate, then measure the absorbance at 450nm using a microplate reader after incubating the plate for 1-4 hours in at 37°C.
Inhibition of MCF-7 cell proliferation after incubation with IL1α for 72h observed by inverted microscope was shown in Figure 1. Cell viability was assessed by CCK-8 (Cell Counting Kit-8) assay after incubation with various concentrations of IL1α for 72h. The mean OD value of MCF-7 assessed by CCK-8 was shown in Figure 2. It was obvious that IL1α significantly decreased cell viability of MCF-7 cells.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.