Active Epidermal Growth Factor (EGF)

Catalog#:PEH12409

概述（Summary）

别称（Alternative Names）	URG; Beta-Urogastrone; Pro-epidermal growth factor
纯度（Purity）	Greater than 90% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	Please contact with the lab for this information
蛋白构建（Construction）	【Asn971-Arg1023】
Accession #	P01133
表达宿主（Host）	E.coli
种属（Species）	Homo sapiens (Human)
预测分子量（Predicted MW）	6.2 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
背景介绍（References）	EGF stimulates the growth of various epidermal and epithelial tissues in vivo and in vitro and of some fibroblasts in cell culture. Magnesiotropic hormone that stimulates magnesium reabsorption in the renal distal convoluted tubule via engagement of EGFR and activation of the magnesium channel TRPM6. Can induce neurite outgrowth in motoneurons of the pond snail Lymnaea stagnalis in vitro.

电泳图（SDS-PAGE image）



Greater than 90% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）



Figure 1. Cell proliferation of Balb/c 3T3 cells after stimulated with rhEGF.
(A) Balb/c 3T3 cultured in DMEM, stimulated with 4 ng/ml EGF 72h;
(B) Unstimulated Balb/c 3T3 cells cultured in 0.4% FBS DMEM for 72h.



Figure 2. The dose-effect curve of recombinant human EGF on Balb/c 3T3 cells

Epidermal growth factor (EGF) is a growth factor that stimulates cell growth, proliferation, and differentiation by binding to its receptor EGFR. To test the effect of EGF on cell proliferation of 3T3 fibroblasts, Balb/c 3T3 cells were seeded into triplicate wells of 96-well plates and allowed to attach overnight, then the medium was replaced with various concentrations of recombinant human EGF. After incubated for 48h, cells were observed by inverted microscope and cell proliferation was measured by Cell Counting Kit-8 (CCK-8). Briefly, 10 µl of CCK-8 solution was added to each well of the plate，then measure the absorbance at 450 nm using a microplate reader after incubating the plate for 1-4 hours at 37 °C. Cell proliferation of Balb/c 3T3 cells after incubation with rhEGF for 48h observed by inverted microscope was shown in Figure 1. The dose-effect curve of rhEGF was shown in Figure2. It was obvious that rhEGF significantly promoted cell proliferation of 3T3 cells. The ED50 for this effect is typically 3.9 ng/ml.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.