
Active Carbonic Anhydrase IX (CA9)

Catalog#:PEH12894

概述（Summary）


别称（Alternative Names）	CA-IX; CAIX; MN; pMW1; P54/58N; Carbonic Dehydratase; Renal cell carcinoma-associated antigen G250; RCC-Associated Protein G250; Carbonate dehydratase IX
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	Please contact with the lab for this information
蛋白构建（Construction）	【Pro59-Asp414】
Accession #	Q16790
表达宿主（Host）	E.coli
种属（Species）	Homo sapiens (Human)
预测分子量（Predicted MW）	42.5 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
背景介绍（References）	Reversible hydration of carbon dioxide. Participates in pH regulation. May be involved in the control of cell proliferation and transformation. Appears to be a novel specific biomarker for a cervical neoplasia.

电泳图（SDS-PAGE image）



Greater than 95% as determined by reducing SDS-PAGE.

活性图（Bioactivity image）



4-Nitrophenol (product)mM	OD400nm
0.625	1.9
0.3125	1.038
0.15625	0.494
0.078125	0.259
0.0390625	0.1
0.01953125	0.066

Figure 1. The standard curve of 4-Nitrophenol

One unit of enzyme activity is defined as the 1 μg of enzyme required to convert 1 pmol of 4-Nitrophenyl acetate to 4-Nitrophenol in 1min at 37°C. The specific activity of recombinant human CA9 is 948 pmol/min/μg.

Specific Activity (pmol/min/μg)= $\frac{\Delta OD * F}{T * N}$

△OD=Adjusted for Substrate Blank
F=Conversion Factor(convert from standard curve of 4-Nitrophenol)
T= Time
N=Amount of enzyme

Carbonic Anhydrase IX (CA9) also known as membrane antigen MN and renal cell carcinoma (RCC) associated antigen G250, is a transmembrane enzyme expressed primarily in carcinoma cells. It is one of the best markers for hypoxia and for RCC. rhCA9 corresponds to the extracellular portion of human CA9. It catalyzes the reversible reaction of CO2 H2O = HCO3- H , which is fundamental to many processes such as respiration, renal tubular acidification and bone resorption. The recombinant human CA9 activity was measured by its ability to hydrolyze 4-Nitrophenyl acetate (4-NPA) to 4-Nitrophenol. The reaction was performed in 12.5 mM Tris, 75 mM NaCl, pH 7.5 (Assay Buffer), initiated by addition 50 μL of various concentrations of CA9 (diluted by Assay Buffer) to 50 μL of 2 mM Substrate 4-NPA (100 mM stock in Acetone, diluted by deionized water). Incubated at 37°C for 5min, then read at a wavelength of 400 nm.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.