Recombinant Diacylglycerol Kinase Zeta (DGKz)(N-6His)


Catalog#:PEH22419

概述（Summary）

别称（Alternative Names）	DAGK5; DAGK6; DGK-ZETA; hDGKzeta; Diglyceride kinase zeta
纯度（Purity）	Greater than 97% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	6His-【Val695-Ala932】
Accession #	Q13574
表达宿主（Host）	E.coli
种属（Species）	Homo sapiens (Human)
预测分子量（Predicted MW）	30.1 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of PBS, pH7.4, containing 0.01% SKL, 1mM DTT, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in 10mM PBS (pH7.4) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
背景介绍（References）	Diacylglycerol kinase that converts diacylglycerol/DAG into phosphatidic acid/phosphatidate/PA and regulates the respective levels of these two bioactive lipids. Thereby, acts as a central switch between the signaling pathways activated by these second messengers with different cellular targets and opposite effects in numerous biological processes. Also plays an important role in the biosynthesis of complex lipids. Does not exhibit an acyl chain-dependent substrate specificity among diacylglycerol species. Can also phosphorylate 1-alkyl-2-acylglycerol in vitro but less efficiently and with a preference for alkylacylglycerols containing an arachidonoyl group. The biological processes it is

involved in include T cell activation since it negatively regulates T-cell receptor signaling which is in part mediated by diacylglycerol. By generating phosphatidic acid, stimulates PIP5KIA activity which regulates actin polymerization. Through the same mechanism could also positively regulate insulin-induced translocation of SLC2A4 to the cell membrane.; Regulates RASGRP1 activity.; Does not regulate RASGRP1 activity.

电泳图（SDS-PAGE image）



Greater than 97% as determined by reducing SDS-PAGE.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.