Recombinant Fucosyltransferase 1 (FUT1)(N-6His)


Catalog#:PEH22602

概述（Summary）

别称（Alternative Names）	H; HH; HSC; Galactoside 2-Alpha-L-Fucosyltransferase,H Blood Group; GDP-L-fucose:beta-D-galactoside 2-alpha-L-fucosyltransferase 1; Blood group H alpha 2-fucosyltransferase
纯度（Purity）	Greater than 95% as determined by reducing SDS-PAGE.
内毒素（Endotoxin level）	<1 EU/μg as determined by LAL test.
蛋白构建（Construction）	6His-【Trp149-Pro365】
Accession #	P19526
表达宿主（Host）	E.coli
种属（Species）	Homo sapiens (Human)
预测分子量（Predicted MW）	28.6 kDa
制剂（Form）	Lyophilized from a 0.2 μm filtered solution of 20mM Tris, 150mM NaCl, pH8.0, containing 1mM EDTA, 1mM DTT, 0.01% SKL, 5% Trehalose and Proclin300.
运输方式（Shipping）	The product is shipped with ice bag. Upon receipt, store it immediately at the temperature listed below.
稳定性&储存（Stability &Storage）	12 months at -20°C to -80°C in lyophilized state.6 months at -20°C to -80°C under sterile conditions after reconstitution.7-10 days at 2°C to 8°C under sterile conditions after reconstitution.Please minimize freeze-thaw cycles.
复溶（Reconstitution）	Reconstitute in 20mM Tris, 150mM NaCl (pH8.0) to a concentration of 0.1-1.0 mg/mL. Do not vortex.
背景介绍（References）	Catalyzes the transfer of L-fucose, from a guanosine diphosphate-beta-L-fucose, to the terminal galactose residue of glycoconjugates through an alpha(1,2) linkage leading to H antigen synthesis that is an intermediate substrate in the synthesis of ABO blood group antigens. H antigen is essential for maturation of the glomerular layer of the main olfactory bulb, in cell migration and early cell-cell contacts during tumor associated angiogenesis. Preferentially fucosylates soluble lactose and to a lesser extent fucosylates glycolipids gangliosides GA1

and GM1a.

电泳图（SDS-PAGE image）



Greater than 95% as determined by reducing SDS-PAGE.

注意事项（Note）

For Research Use Only , Not for Diagnostic Use.